

# Results from BNL E949 on

$$K^+ \rightarrow \pi^+ \gamma \gamma$$

**Benji Lewis**  
**University of New Mexico**

**Tamaki Yoshioka**  
**KEK**

**P. Kitching**

*Centre for Subatomic Research, University of Alberta*

**D.A. Bryman**

*University of British Columbia*

**B. Bhuyan**, I-H. Chiang, M.V. Diwan, J.S. Frank, J.S. Haggerty, D.E. Jaffe, S.H. Kettell, K.K. Li,

L.S. Littenberg, G. Redlinger, R.C. Strand and **B. Viren**

*Brookhaven National Laboratory (BNL)*

**P.S. Cooper**, **E. Ramberg** and **R.S. Tschirhart**

*Fermi National Accelerator Laboratory (FNAL)*

**M. Miyajima** and **Y. Tamagawa**

*Fukui University*

**A. Artamonov**, **A. Kozhevnikov**, **A. Kushnirenko**, L. Landsberg, V. Mukhin, V. Obraztsov, D. Patalakha,

**S. Petrenko** and **D. Vavilov**

*Institute for High Energy Physics (IHEP)*

**V.V. Anisimovsky**, **A.P. Ivashkin**, **M.M. Khabibullin**, **A.N. Khotjantsev**, **Y.G. Kudenko**, **O.V. Mineev** and

**N.V. Yershov**

*Institute for Nuclear Research (INR)*

**S. Kabe**, **M. Kobayashi**, **T.K. Komatsubara**, **E. Ohashi**, **K. Omata**, **T. Sato**, **T. Sekiguchi**, **S. Sugimoto**,

**T. Tsunemi**, **Y. Yoshimura** and **T. Yoshioka**

*High Energy Accelerator Research Organization (KEK)*

**N. Muramatsu**

*Japan Atomic Energy Research Institute (JAERI)*

**T. Fujiwara**, **K. Mizouchi**, **T. Nomura** and **N. Sasao**

*Kyoto University*

**T. Shinkawa**

*National Defense Academy of Japan*

**B. Bassalleck**, **B. Lewis** and **J. Lowe**

*University of New Mexico (UNM)*

**M. Nomachi**

*Osaka University*

**T. Nakano**

*Research Center for Nuclear Physics (RCNP), Osaka University*

**I.-A. Christidi** and **M.D. Marx**

*Stony Brook University*

**P.C. Bergbusch**, **E.W. Blackmore**, **S. Chen**, **J. Hu**, **A. Konaka**, **J.A. Macdonald**, **J. Mildenberger**,


**T. Numao**, **J.-M. Poutissou** and **R. Poutissou**

*TRIUMF*

## $\pi^+$ Momentum Spectrum for E787


Below  $K^+ \rightarrow \pi^+ \pi^0$  peak     $\mathcal{B}(K^+ \rightarrow \pi^+ \gamma\gamma) = [6.0 \pm 1.5 \pm 0.7] \times 10^{-7}$

Above  $K^+ \rightarrow \pi^+ \pi^0$  peak     $\mathcal{B}(K^+ \rightarrow \pi^+ \gamma\gamma) < 5.0 \times 10^{-7}$



P. Kitching *et al.*, Phys. Rev. Lett. **79**, 4079 (1997).

## Branching Ratio from ChPT



- $O(p^6)$  corrections increase BR by 30-40%.
- Finite BR at kinematic end point.

Values obtain from E787 1991


$$\hat{c}_{O(p^4)} = 1.6 \implies \mathcal{B} = 4.9 \times 10^{-10}$$

$$\hat{c}_{O(p^6)} = 1.8 \implies \mathcal{B} = 6.1 \times 10^{-9}$$


Order of Magnitude  
different!

# E949 Detector & Trigger

- $K^+$  enters detector & stops.
- $K^+$  decays : detect  $\pi^+$  and  $\gamma$ s
- Trigger on 1 or 2  $\gamma$  clusters
  - $\gamma$  in Barrel photon detectors
  - $2\gamma \rightarrow 1 \gamma\text{cluster}$  for high mom.  $\pi^+$
- long range  $\pi^+$



# Backgrounds & Trigger Events



E787 Signal  
Region


E949 Signal  
Region

$K^+ \rightarrow \pi^+\pi^0; \pi^0 \rightarrow \gamma\gamma$   
**SAME FINAL STATE!**

Background  
from  $K^+ \rightarrow \pi^+\pi^0$


$K^+ \rightarrow \mu^+\nu\gamma$   
 $K^+ \rightarrow \mu^+\pi^0\nu$

$K^+ \rightarrow \pi^+\gamma\gamma$  triggers




# $K^+ \rightarrow \pi^+ \pi^0$ Background

$K^+ \rightarrow \pi^+ \gamma\gamma$  signal




background from  $K^+ \rightarrow \pi^+ \pi^0$




## Suppress backgrounds

- Use  $\pi^+$  kinematics
- Opening angle cut on  $\gamma$ s



# Overlapping $\gamma$ Background



- *Cut on overlapping  $\gamma$  by observing larger than expected energy*

## Results

Background	Background level
$K^+ \rightarrow \pi^+ \pi^0$	$0.017 \pm 0.006$
<i>Overlapping <math>\gamma</math></i>	$0.065 \pm 0.065$
Muon	$0.090 \pm 0.020$
Single Beam	$0.025 \pm 0.014$
Double Beam	$0.006 \text{ (90\% } C.L.)$
<b>Total</b>	$0.197 \pm 0.070$

## Acceptance


$$A_{O(p^6)}^{\pi^+ \gamma \gamma} = 1.550 \pm 0.034 \times 10^{-4}$$

## Expected Number of Events


1.6

# Open Box

$K^+ \rightarrow \pi^+ \gamma\gamma$   
Signal Region




Surviving events  
All cuts applied



Simulated distribution  
 $\hat{c} = 1.8$  unitarity corrections

# Conclusions


$$\mathcal{B}(K^+ \rightarrow \pi^+ \gamma\gamma; P_{\pi^+} > 213 \text{ MeV}/c) < 9.1 \times 10^{-9} \quad (90\% \text{ CL})$$



- Upper limit 7.6x improvement
- consistent with ChPT  $O(p^6)$  &  $O(p^4)$
- More data needed.


THANK YOU

- Upper



**Extras**

$$K^+ \rightarrow \pi^+ X^0, X^0 \rightarrow \gamma\gamma$$



## E949 Method

$K^+$  enters E949 Detector



- Cerenkov:  $K^+/\pi^+$  (PID)
- Active Degradar: slows  $K^+$
- Target:  $K^+$  comes to rest.
- Beam: no other particles (PID)

$K^+$  decays

- Target: 'Delayed Coincidence' between  $K^+$  and  $\pi^+$  (Timing)
- Photo Veto: extra E (Veto)
- Target: E,R,T (Kinematics)
- Drift Chamber: P (Kinematics)
- Range Stack: E,R (Kinematics)

$\pi^+$  decays




- Range Stack:  
 $\pi^+ \rightarrow \mu^+ \rightarrow e^+$  (PID)




# Reconstructing Photons in BV & BVL

- *Trigger on 1 or 2  $\gamma$  clusters*
- *$2 \gamma \rightarrow 1 \gamma \text{cluster}$  for high mom.  $\pi^+$*

## Signature of Signal




# Side view of E949 detector





# Differential Branching Ratio

*ChPT with  $O(p^4)$*





$\hat{c} = -2$  (solid),  $\hat{c} = 0$  (dashed),  $\hat{c} = 2$  (dot – dashed)

# $\pi^+$ Momentum Spectrum



# Branching Ratio from ChPT



Values obtain from E787 1991

$$\hat{C}_{O(p^4)} = 1.6 \implies \mathcal{B} = 4.9 \times 10^{-10}$$

$$\hat{C}_{O(p^6)} = 1.8 \implies \mathcal{B} = 6.1 \times 10^{-9}$$


Order of Magnitude  
different!

# Chiral Perturbation Theory (ChPT)



ChPT prediction for  
 $K^+ \rightarrow \pi^+ \gamma \gamma$

- no tree level contribution  $\mathcal{O}(p^2)$
- $K^+ \rightarrow \pi^+ \gamma \gamma$  amplitude is determined by  $\hat{c}$
- $\hat{c} \equiv \frac{128\pi^2}{3} [3(L_9 + L_{10}) + L_{14} - L_{15} - 2L_{18}]$

loop diagrams from ChPT  
 for  $\mathcal{O}(p^4)$



# Backgrounds & Trigger Events



Backgrounds	Suppression Methods			
	Kinematics	Photon	Particle ID	Timing
$K^+ \rightarrow \pi^+ \pi^0$	✓	✓		
Overlapping Photon	✓		✓	
Muon	✓		✓	
Single Beam			✓	✓
Double Beam				✓ ✓

Backgrounds	Suppression Methods			
	Kinematics	Photon	Particle ID	Timing
$K^+ \rightarrow \pi^+ \pi^0$	✓	✓		
Overlapping Photon	✓		✓	
Muon	✓		✓	
Single Beam			✓	✓
Double Beam				✓ ✓